DRAFT VERSION APRIL 24, 2013
Preprint typeset using LaTeX style emulateapj v. 11/10/09

arXiv:1304.6142v1 [astro-ph.CO] 23 Apr 2013

THE ULTIMATE FATE OF COSMIC RAYS FROM GALAXIES AND THEIR ROLE IN THE INTERGALACTIC MEDIUM

BRIAN C. LACKI[1,2]

Draft version April 24, 2013

ABSTRACT

The majority of cosmic rays (CRs) generated by star-forming galaxies escape those galaxies and enter the intergalactic medium (IGM). I show that the mean pressure of the intergalactic CRs is likely within an order of magnitude of the mean Lyman α forest thermal pressure. At $z \gtrsim 1$, their pressure may have even been dominant. I also demonstrate that, whichever IGM phase the CRs end up in, they are a significant contributor to its pressure support if its temperature is $\sim 10^4$ K, regardless of density. Which phase the CRs end up in depends on the structure and strength of intergalactic magnetic fields. I argue that CRs end up at least 100 kpc from their progenitor galaxies. CRs may self-confine in the IGM to the sound speed, generating $\gtrsim 10^{-13}$ G magnetic fields. These considerations imply the existence and importance of a nonthermal IGM.

Subject headings: intergalactic medium — cosmic rays — galaxies: halos

1. INTRODUCTION

Most of the baryonic matter in the Universe is not in stars or galaxies, but in the gas between the galaxies, the intergalactic medium (IGM). Far from being a uniform background gas, the IGM has a rich structure with several phases spanning a wide range of physical conditions. About one-third of the baryonic mass is in the Lyman α (Lyα) forest, a volume-filling photoionized phase with overdensities $\delta \lesssim 10$ and temperatures 10^4 – 10^5 K (Bi & Davidsen 1997). Before $z \approx 1$, this phase made up most of the baryonic mass of the Universe. As some of this gas collapses into smaller structures, it is shock heated to form the Warm Hot Intergalactic Medium (WHIM), a moderately dense ($\delta \sim 10-100$), 10^5 – 10^7 K phase that presently contains $\sim 1/2$ of the baryonic mass (Cen & Ostriker 1999). Finally, some of the material has collapsed even further into "condensed" structures: these include the galaxies themselves, but also clusters and their hot intracluster media, as well as the multiphased gaseous haloes surrounding galaxies. The circumgalactic gas results from the interaction of galaxies and the IGM, through accretion onto galaxies and expulsion from winds, and itself is complex, with cool dense and warm rarefied phases (e.g., Chen et al. 2001).

But is the picture of the IGM described above, an interplay of thermal gas, galaxies, and gravity, complete? Just as in galaxies, the thermal gas phases described above may be complemented by pervasive nonthermal fields: the cosmic rays (CRs) and magnetic fields. For the special case of galaxy clusters, detections of radio synchrotron emission conclusively demonstrate the presence of both cosmic rays and magnetic fields, and there has been much work seeking an understanding of how these phases are generated and interact with the intracluster gas (e.g., Subramanian et al. 2006; Ferrari et al. 2008). Radio emission is also detected from galaxy filaments, suggesting similar processes are at work (Brown & Rudnick 2009; see also Ryu et al. 2008). On a broader scale, evidence for nonthermal processes in the IGM is scant, but there are theoretical reasons to expect they exist. A nonthermal IGM can take the form of CRs accelerated by structure formation shocks (Loeb & Waxman 2000; Keshet et al. 2004) or injected by dark matter annihilation throughout the Universe (Mapelli et al. 2006), pervasive magnetic fields seeded by early galaxies or even primordially in the Big Bang (Widrow 2002), and through the heating of the IGM by absorption of low frequency nonthermal radio waves (Lacki 2010) or rapid stopping of cascades generated by high energy gamma rays (Broderick et al. 2012; Chang et al. 2012; for the debate on whether this actually happens, see Miniati & Elyiv 2012; Schlickeiser et al. 2012).

A guaranteed source of nonthermal energy in the IGM is the CRs accelerated by star-forming galaxies. In galaxies like the present-day Milky Way, most of the CR energy is in protons. They diffuse out of the galaxy, with only a few percent of the energy lost in pion production and ionization (Strong et al. 2010). Starbursts and higher redshift normal galaxies have more gas, with stronger energy losses. However, γ-ray observations of the starbursts M82 and NGC 253 indicate that $\sim 20-40\%$ of their CR power is lost to pion production, with the majority of that power ($\sim 60-80\%$) escaping the galaxy, most likely in a starburst wind (Lacki et al. 2011; Abramowski et al. 2012; Ackermann et al. 2012). Thus, a large fraction of the power injected in CRs in the Universe ends up in the IGM. Over the Gyrs of cosmic star formation, a significant intergalactic cosmic ray population grew.

Yet the effects of this population has been largely ignored. Nath & Biermann (1993) and Samui et al. (2005) have studied how CRs from star-forming galaxies could have reionized the IGM, or heated it at high redshift. In general, though, the energy losses for typical CRs with kinetic energies of GeV or more are negligible in the present IGM. The pionic loss lifetime for CR protons in the $z = 0$ IGM is $t_\pi \approx 2 \times 10^{14} (1+\delta)^{-1}$ yr for an intergalactic density $n_H = 2.5(1+\delta) \times 10^{-7}\ \mathrm{cm}^{-3}$ (Mannheim & Schlickeiser 1994). The ionization lifetime is even greater for particles with kinetic energies greater than a few hundred MeV. A few other estimates of the intergalactic CR spectrum, with widely disparate assumptions, are briefly given in Dar & de Rújula (2005) and Lipari & et al. (2005).

CRs can nevertheless couple dynamically to the IGM and its weak magnetic fields. I show in this Letter that the mean intergalactic energy density of CRs from star-forming galaxies is probably within a factor of a few of the Lyα forest pressure (Section 2). I also show that CRs are likely to reach the IGM, where they excite magnetic fluctuations in the IGM as

---

[1] Jansky Fellow of the National Radio Astronomy Observatory
[2] Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA, brianlacki@ias.edu

they attempt to self-confine (Section 3). I assume $\Omega_\Lambda = 0.75$, $\Omega_m = 0.25$, $H_0 = 70$ km s^{-1} Mpc^{-1} for the cosmology. For the baryonic density, I assume $\Omega_b = 0.045$ and mean molecular weight $\mu = 0.6$ for a mean IGM comoving number density $\langle n_{\rm IGM} \rangle = 4.1 \times 10^{-7}$ cm^{-3}.

2. THE PRESSURE OF INTERGALACTIC COSMIC RAYS

CR nuclei experience only adiabatic losses once they leave a galaxy. First, CRs in a galactic wind lose momentum adiabatically as the wind expands, sacrificing their energy to push the wind outwards (Völk et al. 1996). Galactic winds are common in galaxies with high specific star-formation rate, both high-z main sequence galaxies and compact starbursts (Heckman et al. 1990; Steidel et al. 2010). Those winds are powered by the efficient thermalization of supernova mechanical energy. Galaxies without supernova-launched winds may still have more rarefied winds originating from the halo and powered by CRs, as CR streaming excite plasma waves that push halo plasma out (Breitschwerdt et al. 1991). Second, CRs in the IGM lose momentum as the Universe expands.

Adiabatic losses conserve the number of particles per logarithmic bin in momentum. The equation for the IGM particle spectrum at redshift z is therefore

$$\frac{dN}{d\ln p}(z) = \int_z^\infty \frac{dQ}{d\ln p_{\rm inj}}(p_{\rm inj}, z_{\rm inj}) \frac{dt}{dz_{\rm inj}} dz_{\rm inj}. \qquad (1)$$

Particles are accelerated to momenta at $p_{\rm inj}$ at a rate $dQ/d\ln p_{\rm inj}$ at redshift $z_{\rm inj}$. I relate $p_{\rm inj}$ to p as $p(z) = \epsilon_{\rm adv} p_{\rm inj}(z_{\rm inj})(1+z)/(1+z_{\rm inj})$, where $\epsilon_{\rm adv}$ accounts for adiabatic losses within a galactic wind. I ignore ionization losses.

CRs advected out from their host galaxy in a wind with density ρ lose momentum as $\rho^{-1/3}$, so the $\epsilon_{\rm adv}$ factor is simply the cube root of the ratio of the final and initial wind density. For example, from the Milky Way, the wind initially occupies the disk with radius 10 kpc and the CRs move with the Alfven speed 10 km s^{-1}. If the wind outflows spherically and reaches 1 Mpc with a speed of 300 km s^{-1} (Breitschwerdt et al. 1991), then $\epsilon_{\rm adv} = [(4\pi(1~{\rm Mpc})^2(300~{\rm km~s^{-1}}))/(2\pi(10~{\rm kpc})^2(10~{\rm km~s^{-1}}))]^{-1/3} \approx 0.012$. Adiabatic cooling is a much greater obstacle within winds from small, compact starburst regions; the original volume is small and the termination shocks are generally far from the burst, resulting in a huge density contrast.

Although winds could drain most of the CR energy by the time it reaches the IGM, the wind itself eventually stops at a termination shock when its ram pressure equals the surrounding IGM's pressure. The termination shock itself can accelerate CRs, converting $\epsilon_{\rm shock} \gtrsim 10\%$ of the wind's kinetic energy back into CRs and releasing them into the IGM (Jokipii & Morfill 1985). For Milky Way-like galaxies, the power injected at these shocks is $\epsilon_{\rm shock}$ times the luminosity in CRs in the galaxy, whereas in starburst shocks, the power injected is $\epsilon_{\rm shock}$ times the supernova mechanical power, about 10 times greater per unit star formation. Reacceleration can also occur at shocks within inhomogeneous winds (Dorfi & Breitschwerdt 2012).

The injection spectrum is a power law in momentum ($d^2Q/(dp_{\rm inj}~dt) = Cp_{\rm inj}^{-\Gamma}$), with spectral index $\Gamma = 2.2$. The normalization of the spectrum is set by a volumetric energy injection rate:

$$\dot{\varepsilon}_{\rm CR} = \int_{{\rm MeV}/c}^{{\rm PeV}/c} CK p_{\rm inj}^{-\Gamma} dp_{\rm inj}, \qquad (2)$$

where K is kinetic energy. I calculate $\dot{\varepsilon}_{\rm CR}$ by scaling to the cosmic star-formation rate density (Hopkins & Beacom 2006). Star formation creates young, massive stars which accelerate the CRs. The star-formation rate is directly related to the supernova rate as $\Gamma_{\rm SN} = 0.0084~{\rm yr}^{-1}({\rm SFR}/(M_\odot~{\rm yr}^{-1}))$. Multiwavelength models of star-forming galaxies indicate that each supernova accelerates $\sim 10^{50}$ erg of CRs (that is, 10% of the 10^{51} erg released in mechanical energy). So I have

$$\dot{\varepsilon}_{\rm CR} = 0.018~{\rm meV~Gyr^{-1}~cm^{-3}} \left(\frac{\rho_{\rm SFR}}{1~M_\odot~{\rm yr^{-1}~Mpc^{-3}}}\right). \qquad (3)$$



FIG. 1.— Predicted comoving cosmic CR energy density. The dash-dotted line is if 10% of the original power in CRs is accelerated at termination shocks, while solid lines represent the original population of CRs after experiencing adiabatic losses. The grey shaded band is the expected mean IGM thermal pressure for $T = 10^4$ K – 10^6 K.

With equations 1–3, I calculate the evolution of the intergalactic CR energy density $U_{\rm CR}$ with redshift. The solid lines in Figure 1 are the expected $U_{\rm CR}$ of the original CR population with $\epsilon_{\rm adv}$ =0.001 – 1. In addition, I plot the $U_{\rm CR}$ from CRs accelerated at galactic wind termination shocks with $\epsilon_{\rm shock} = 0.1$ of the original CR power converted back into CRs (dash-dotted line).

I find that with $\epsilon_{\rm adv} = 1$, the comoving CR energy density $U'_{\rm CR}(z)/k_B$ is 0.02 K cm^{-3} at present, and it peaked at $z = 1.0$ at 0.1 K cm^{-3}. For termination shocks CRs with $\epsilon_{\rm shock} = 0.1$, $U'_{\rm CR}/k_B$ is ten times smaller. For comparison, the expected comoving thermal IGM energy densities are 0.0075 – 0.75 K cm^{-3} for $T = 10^4$–10^6 K (grey band in Figure 1). We see that, if $\epsilon_{\rm adv} = 1$, intergalactic CRs have an energy density greater than that of the Lyα forest. Even for a conservative $\epsilon_{\rm shock} = 0.1$, $U_{\rm CR}$ is roughly equal to the Lyα forest thermal energy density at $z \approx 1$ and are a quarter of it at $z \approx 0$. Thus CRs may play an important role in the dynamics of the IGM.

2.1. *CR importance in other IGM phases*

CRs might not actually fill the entire volume of the Universe, but instead may be confined to one IGM phase. These phases can be very overdense. But then since the same CR power is being squeezed into a smaller confinement volume, the CR energy density is likewise greater.

What is the ratio of CR energy density to thermal energy density in the phase the CRs end up in? Suppose the mean CR energy density of the Universe is $\langle U_{\rm CR}\rangle$. If a fraction f_i of the CR energy occupies an IGM phase i, and if that phase has a cosmic filling factor ϕ, then the CR energy density within that phase is:

$$U_{\rm CR} = f_i \phi \langle U_{\rm CR}\rangle. \tag{4}$$

The gas density in that phase is limited by mass conservation to:

$$n_i \le \frac{\Omega_i}{\Omega_b}\phi\langle n_{\rm IGM}\rangle, \tag{5}$$

where Ω_i is the cosmic volume-averaged fraction of the critical density that the IGM phase accounts for. The thermal energy density within that phase is $U_i = (3/2)n_i k T_i$, giving the ratio

$$\frac{U_{\rm CR}}{U_{\rm IGM}} = \frac{\Omega_b}{\Omega_i} f_i \frac{2\langle U_{\rm CR}\rangle}{3 n_i k T_i}. \tag{6}$$

The filling factors cancel out: if CRs end up mostly in one phase of the IGM, then the only relevant IGM parameters are its mass fraction (which is always less than 1) and its temperature. The CR energy density exceeds the thermal energy density as long as:

$$T_i < \frac{\Omega_b}{\Omega_i} f_i \frac{2\langle U_{\rm CR}\rangle}{3\langle n_{\rm IGM}\rangle k}. \tag{7}$$

At $z = 0$, the CRs dominate the energy density of their final phase if

$$T_i < 3.6\times 10^4 {\rm K}\frac{\Omega_b}{\Omega_i} f_i \Phi, \tag{8}$$

where $\Phi \approx 0.1$ is the ratio of the actual $U_{\rm CR}^{\prime}$ and its value for $\epsilon_{\rm adv} = 1$. About half of the $z = 0$ IGM mass has a temperature $\sim 10^4$ K, consisting of the Lyman α forest and condensed haloes, indicating a major CR contribution to the pressure. However, half of the IGM mass is WHIM with $10^5 - 10^7$ K temperatures; if the CRs end up in WHIM, they are insignificant.

At higher redshift, CRs are even more important because $\rho_{\rm SFR}$ was much greater. Furthermore, an increasingly larger mass fraction of the gas is in the uncollapsed 10^4 K Lyα forest. At $z = 1$, I find that CRs dominate the pressure of their final phase if

$$T_i(z = 1) < 1.6\times 10^5\ {\rm K}\frac{\Omega_b}{\Omega_i} f_i \Phi, \tag{9}$$

a condition which held for most of the IGM's mass.

Thus, CRs are likely to provide significant pressure support in some IGM phase at $z \gtrsim 1$.

3. WHERE ARE THE COSMIC RAYS?

3.1. *Do CRs really even escape their galaxies?*

Observations of quasar absorption lines indicate that winds in high-z star-forming galaxies transport material out to $\gtrsim$ 100 kpc (Heckman et al. 1990). Theory likewise indicates that even Milky Way galaxies host CR-driven winds (Breitschwerdt et al. 1991). Ultimately, a wind with asymptotic speed v_∞ and ram luminosity $\dot{E}$ should flow out to the termination shock radius $R_s = \sqrt{\dot{E}/(4\pi v_\infty P_{\rm IGM})}$, or

$$R_s \approx 1.1\ {\rm Mpc}\left(\frac{\dot{E}}{10^{40}\ {\rm erg\ sec^{-1}}}\right)^{1/2}\left(\frac{v_\infty}{1000\ {\rm km\ s^{-1}}}\right)^{-1/2} \times\left(\frac{P_{\rm IGM}/k_B}{0.005\ {\rm K\ cm^{-3}}}\right)^{-1/2}. \tag{10}$$

It takes roughly 1 (10) Gyr for a 1000 (100) km s^{-1} wind to traverse 1 Mpc. Thus, galactic winds can transport energy well into the IGM.

But suppose CRs are not advected out but simply diffuse from their host galaxies? CR diffusion results in a net flow out of the Galaxy only if the energy density outside is less than that inside. Since CRs do actually diffuse out of the Galaxy, the CR energy density in the Galactic halo is less than that within the disk. For a steady CR luminosity, we therefore have

$$\frac{t_{\rm MW}}{V_{\rm out}} < \frac{t_{\rm in}}{V_{\rm in}}. \tag{11}$$

The time that the Milky Way has been forming stars, $t_{\rm MW}$, is about 10 Gyr, whereas the time that CRs stay within the Galactic disk is only about $t_{\rm in} \approx$30 Myr (Connell 1998). The confinement volume of the Galactic disk is $\gtrsim$ 350 kpc^3 (Schlickeiser 2002). Supposing that the CRs fill a sphere with radius $R_{\rm out}$, the diffusion of CRs out of the Milky Way conservatively gives us:

$$R_{\rm out} \gtrsim \left(\frac{3}{4\pi}V_{\rm in}\frac{t_{\rm MW}}{t_{\rm in}}\right)^{1/3} \gtrsim 30\ {\rm kpc}\left(\frac{V_{\rm in}}{350\ {\rm kpc}^3}\right)^{1/3}. \tag{12}$$

If CRs are not advected away, they *must* diffuse far beyond the star-formation and gas in the Galactic disk, well into the circumgalactic gas.

3.2. *How far in the IGM do they go?*

The propagation of CRs depends on the magnetic field strength and structure in the IGM (Adams et al. 1997). Very little is known about the magnetic field strength, other than that it less than $\lesssim 1\ \mu$G on all coherence lengths and $\lesssim$ 1 nG on large scales (Neronov & Vovk 2010). The most interesting constraints come from gamma-ray observations. The lack of GeV cascade emission from blazars suggests that pair $e^\pm$ generated by TeV γ-rays on their way to Earth are deflected out of the sightline to Earth by intergalactic magnetic fields (e.g., Neronov & Vovk 2010). This sets a lower limit of 10^{-18} G (Dermer et al. 2011), but the applicability of the limits has been disputed (Broderick et al. 2012).

Supposing that CRs do reach the large-scale IGM, the slowest they can diffuse is Bohm diffusion – one scattering per gyroradius. With a mean free path of the gyroradius,

$$r_L = 1.1\ {\rm Mpc}\ \left(\frac{E}{\rm GeV}\right)\left(\frac{B}{10^{-18}\ {\rm G}}\right)^{-1}, \tag{13}$$

the distance CRs can diffuse is $s_{\rm Bohm} = \sqrt{ctr_L}$

$$s_{\rm Bohm} = 58\ {\rm Mpc}\ \left(\frac{t}{10\ {\rm Gyr}}\right)^{1/2}\left(\frac{E}{\rm GeV}\right)^{1/2}\left(\frac{B}{10^{-18}\ {\rm G}}\right)^{-1/2}. \tag{14}$$

The maximum distance that CRs propagate into the IGM is set by the plasma waves that they excite while they stream. Within galaxies, where the magnetic energy density is comparable to thermal pressure, CRs self-confine themselves to speeds less than the Alfven speed (Kulsrud & Pearce 1969). In the classical version of CR self-confinement, the magnetic fluctuations from CR streaming are much smaller than the mean field. However, it is thought that some nonlinear version of the instability can amplify magnetic fluctuations until they are larger than the mean field to confine CRs if necessary (Lucek & Bell 2000).

The sound speed is much greater than the Alfven speed in the IGM, though. In these conditions, the streaming speed of CRs is probably of order the sound speed $c_s = 15 \text{ km s}^{-1} \sqrt{T / 10^4 \text{ K}}$ (Holman et al. 1979). This weak self-confinement limits the distance that intergalactic CRs travel to:

$$s_{\rm max} = c_s t = 0.15 \text{ Mpc} \left(\frac{T}{10^4 \text{ K}}\right)^{1/2} \left(\frac{t}{10 \text{ Gyr}}\right). \qquad (15)$$

Note that in hot phases like the WHIM, where the CR pressure is least significant, the CRs can stream out fastest. For $T = 10^6$ K, CRs can traverse about 2 Mpc, bigger than the typical size of a WHIM structure. Streaming CRs may escape quickest from those structures where they are least important.

Even if wave generation slows down CRs in the IGM, this is very interesting in itself: it means that CRs excite magnetic fields in the IGM. The CR mean free path is at least a Larmor radius. If their average streaming speed over a time $t_{\rm stream}$ is $\lesssim c_S$, then the magnetic field must be at least $B \gtrsim cE / (e c_s^2 t_{\rm stream})$, where E is the particle energy and e is the particle charge. The characteristic magnetic field strengths are

$$B \gtrsim 10^{-13} \text{ G} \left(\frac{E}{\text{GeV}}\right) \left(\frac{T}{10^4 \text{ K}}\right)^{-1} \left(\frac{t_{\rm stream}}{10 \text{ Gyr}}\right)^{-1} \qquad (16)$$

on some coherence length greater than $r_L = 10 \text{ pc } (E/\text{GeV})(B/10^{-13} \text{ G})^{-1}$. The existence of these fluctuations is consistent with current limits on intergalactic magnetic fields (Neronov & Vovk 2010). The magnetic fluctuations that CRs can excite can have no larger than $\sim \sqrt{8\pi U_{\rm CR}}$ on energetics grounds; for $z = 0$, this sets an upper limit of 10 nG.

4. CONCLUSIONS

The injection of CRs by star-forming galaxies into the IGM is a long-neglected kind of feedback. The energy in CRs accelerated by star-forming galaxies over the Universe's lifetime is comparable to the thermal energy in the Lyα forest. Although the CRs may experience strong adiabatic losses as they are advected away from their galaxies by winds, some of that energy is recovered in CRs when the winds reach their termination shocks. The calculated pressure of intergalactic CRs is therefore still within an order of magnitude of that of the Lyα forest. If the CRs are trapped in a denser phase, their own density is also necessarily higher, meaning that these CRs are important if they end up in gas with $T \approx 10^4$ K of any density. CRs may be especially important at $z \gtrsim 1$, when the star-formation rate density was greatest and when most of the IGM was relatively cool.

If standard theories of CR streaming apply to the IGM, the CRs affect its magnetic structure. As they self-confine to roughly the sound speed, they excite magnetic fluctuations with $B \gtrsim 10^{-13}$ G. Even so, they can penetrate through up to 100 kpc of 10^4 K gas and 1 Mpc of 10^6 K gas.

Star-forming galaxies are the not the only source of cosmic rays in the IGM. Active galactic nuclei and structure formation contribute CRs too. Rather, CRs from star-forming galaxies demonstrate the need to consider nonthermal processes in the IGM. Indeed, the importance of the nonthermal IGM is beginning to be recognized in other contexts. The nonthermal fields may prove to be just as important to the IGM as in the ISM.

I was supported by a Jansky Fellowship from the National Radio Astronomy Observatory during this work. The National Radio Astronomy Observatory is operated by Associated Universities, Inc., under cooperative agreement with the National Science Foundation.

REFERENCES

Abramowski, A., Acero, F., Aharonian, F., et al. 2012, ApJ, 757, 158
Ackermann, M., Ajello, M., Allafort, A., et al. 2012, ApJ, 755, 164
Adams, F. C., Freese, K., Laughlin, G., Schwadron, N., & Tarle, G. 1997, ApJ, 491, 6
Bi, H., & Davidsen, A. F. 1997, ApJ, 479, 523
Breitschwerdt, D., McKenzie, J. F., & Völk, H. J. 1991, A&A, 245, 79
Broderick, A. E., Chang, P., & Pfrommer, C. 2012, ApJ, 752, 22
Brown, S., & Rudnick, L. 2009, AJ, 137, 3158
Cen, R., & Ostriker, J. P. 1999, ApJ, 514, 1
Chang, P., Broderick, A. E., & Pfrommer, C. 2012, ApJ, 752, 23
Chen, H.-W., Lanzetta, K. M., & Webb, J. K. 2001, ApJ, 556, 158
Connell, J. J. 1998, ApJ, 501, L59
Dar, A., & de Rújula, A. 2005, Phys. Rev. D, 72, 123002
Dermer, C. D., Cavadini, M., Razzaque, S., et al. 2011, ApJ, 733, L21
Dorfi, E. A., & Breitschwerdt, D. 2012, A&A, 540, A77
Ferrari, C., Govoni, F., Schindler, S., Bykov, A. M., & Rephaeli, Y. 2008, Space Sci. Rev., 134, 93
Heckman, T. M., Armus, L., & Miley, G. K. 1990, ApJS, 74, 833
Holman, G. D., Ionson, J. A., & Scott, J. S. 1979, ApJ, 228, 576
Hopkins, A. M., & Beacom, J. F. 2006, ApJ, 651, 142
Jokipii, J. R., & Morfill, G. E. 1985, ApJ, 290, L1
Jokipii, J. R., & Morfill, G. 1987, ApJ, 312, 170
Keshet, U., Waxman, E., & Loeb, A. 2004, ApJ, 617, 281
Kulsrud, R., Pearce, W. P. 1969, ApJ 156, 445
Lacki, B. C. 2010, MNRAS, 406, 863
Lacki, B. C., Thompson, T. A., Quataert, E., Loeb, A., & Waxman, E. 2011, ApJ, 734, 107
Lipari, P., & et al. 2005, International Cosmic Ray Conference, 6, 381
Loeb, A., & Waxman, E. 2000, Nature, 405, 156
Lucek, S. G., & Bell, A. R. 2000, MNRAS, 314, 65
Mannheim, K., & Schlickeiser, R. 1994, A&A, 286, 983
Mapelli, M., Ferrara, A., & Pierpaoli, E. 2006, MNRAS, 369, 1719
Miniati, F., & Elyiv, A. 2012, arXiv:1208.1761
Nath, B. B., & Biermann, P. L. 1993, MNRAS, 265, 241
Neronov, A., & Vovk, I. 2010, Science, 328, 73
Ryu, D., Kang, H., Cho, J., & Das, S. 2008, Science, 320, 909
Samui, S., Subramanian, K., & Srianand, R. 2005, International Cosmic Ray Conference, 9, 215
Schlickeiser, R. 2002, *Cosmic Ray Astrophysics*, (New York: Springer)
Schlickeiser, R., Ibscher, D., & Supsar, M. 2012, ApJ, 758, 102
Steidel, C. C., Erb, D. K., Shapley, A. E., et al. 2010, ApJ, 717, 289
Strong, A. W., Porter, T. A., Digel, S. W., Jóhannesson, G., Martin, P., Moskalenko, I. V., Murphy, E. J., & Orlando, E. 2010, ApJ, 722, L58
Subramanian, K., Shukurov, A., & Haugen, N. E. L. 2006, MNRAS, 366, 1437
Völk, H. J., Aharonian, F. A., & Breitschwerdt, D. 1996, Space Sci. Rev., 75, 279
Widrow, L. M. 2002, Reviews of Modern Physics, 74, 775